



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/55, 31/495, 31/445, 31/16</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/35584</b> <b>(43) International Publication Date:</b> 2 October 1997 (02.10.97)
<b>(21) International Application Number:</b> PCT/US97/04699 <b>(22) International Filing Date:</b> 24 March 1997 (24.03.97) <b>(30) Priority Data:</b> 60/014,152 25 March 1996 (25.03.96) US <b>(71) Applicant:</b> ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US). <b>(72) Inventors:</b> HELTON, David, R.; 2607 South Bo-Mar Lane, Greenfield, IN 46140 (US). SHANNON, Haarlán, E.; 4229 Rolling Springs Drive, Carmel, IN 46234 (US). WOMER, Daniel, E.; Apartment H., 2121 Blue Jay Court, Indianapolis, IN 46260 (US). <b>(74) Agents:</b> VORNDRAN-JONES, MaCharri et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHOD FOR TREATING PAIN  <b>(57) Abstract</b>  The present invention provides a method for treating pain using an atypical antipsychotic compound.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## METHOD FOR TREATING PAIN

5 This invention provides a method for using an atypical antipsychotic compound selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine for the treatment of pain.

10 This invention relates to the treatment of pain using atypical antipsychotic compounds to provide analgesic activity.

Surprisingly, we have discovered that atypical antipsychotic compounds can be particular useful for treating pain. The analgesic effect may be further enhanced when used in combination with one or more another Drug Used in the Treatment of Pain compounds. More specifically, the invention provides a method of treating pain in humans using an atypical antipsychotic compound.

20 There are drugs used in the treatment of pain which known in the literature and to the skilled artisan. see for example, Merck Manual, 16th Ed. (1992) p. 1409.

More active analgesics are in constant demand because they offer the attractive possibility of relieving pain with reduced dosages, thereby diminishing the expected side effects and toxicity that would otherwise result from higher dosages. It would be particularly desirable to acquire a synergistic combination effect to further reduce dosages and diminish side effects. Such a composition is a subject of the present invention.

30 Certain compounds have been disclosed as being atypical antipsychotics which can be useful for treating schizophrenia or related psychotic conditions. Applicants have discovered that atypical antipsychotic compounds selected from the group consisting of risperidone, 35 clozapine, seroquel, sertindole, ziprasidone, and zotepine can be useful for the treatment of pain and may provide a

synergistic effect when administered with one or more other drugs used in the treatment of pain.

5 The present invention provides a method for treating pain, comprising administering an effective amount of an atypical antipsychotic selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine to a patient in need thereof.

10 The present invention further provides a method for treating pain comprising administering to a patient in need thereof, an analgesic composition comprising an atypical antipsychotic or a pharmaceutically acceptable salt thereof; and another Drug Used in the Treatment of Pain, in a weight ratio of one part atypical antipsychotic to from  
15 about one part to about one thousand (1,000) parts of another Drug Used in the Treatment of Pain.

A preferred composition is a weight ratio of atypical antipsychotic to another Drug Used in the Treatment of Pain of from about 1 part atypical  
20 antipsychotic to from about 1 part to about 100 parts of another Drug Used in the Treatment of Pain. An especially preferred ratio is from about 1 part atypical antipsychotic to from about 1 to about 30 parts another Drug Used in the Treatment of Pain. A further preferred  
25 ratio may be from about 1 part atypical antipsychotic to from about 1 part to about 10 parts another Drug Used in the Treatment of Pain. A final preferred ratio may be from about 1 part atypical antipsychotic to from about 1 to about 3 parts another Drug Used in the Treatment of  
30 Pain.

Preferably another Drug Used in the Treatment of Pain is one or more compounds selected from the group consisting of aspirin, acetaminophen, paracetamol, indomethacin, Tylenol #3, tricyclic antidepressants (for  
35 example desipramine, imipramine, amitriptyline, nortriptyline), anticonvulsants (for example, carbamazepine, valproate), and serotonin reuptake

inhibitors (for example, fluoxetine, paroxetine, sertraline), mixed serotonin-norepinephrine reuptake inhibitors (for example venlafaxine, duloxetine), serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, adrenergic agents, and neurokinin antagonists.

Particularly preferred Drug Used in the Treatment of Pain are selected from the group consisting of aspirin, acetaminophen, ketorolac, allopurinol, methysergide maleate, and methotrimeprazine.

The invention further provides a composition for treating pain comprising an atypical antipsychotic or a pharmaceutically acceptable salt or solvate thereof and one or more another Drug Used in the Treatment of Pain in a weight ratio of atypical antipsychotic to another Drug Used in the Treatment of Pain of from about one (1) part atypical antipsychotic to from about 1 part to about 1000 parts Drug Used in the Treatment of Pain.

Another Drug Used in the Treatment of Pain used primarily for the symptomatic relief of pain may be divided into four major groups: 1) opiate analgesics; 2) nonopiate analgesics; 3) analgesics and antipyretics; and 4) nonsteroidal antiinflammatory drugs. Other compounds contemplated herein as "Drug Used in the Treatment of Pain" include, but are in no way limited to other drug classes which might be used with atypical antipsychotics for the treatment of pain to provide a synergistic effect, for example, acetaminophen, paracetamol, indomethacin, Tylenol #3, tricyclic antidepressants (for example desipramine, imipramine, amitriptyline, nortriptyline), anticonvulsants (for example, carbamazepine, valproate), and serotonin reuptake inhibitors (for example, fluoxetine, paroxetine, sertraline), mixed serotonin-norepinephrine reuptake inhibitors (for example venlafaxine, duloxetine), serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, adrenergic agents, and neurokinin

-4-

antagonists. Some preferred another Drug Used in the Treatment of Pain s are selected from acetaminophen, cholinergic analgesics, and neurokinin antagonists. Other preferred Drug Used in the Treatment of Pain include

5 tricyclic antidepressants, anticonvulsants, and serotonin reuptake inhibitors.

Another preferred group of Drug Used in the Treatment of Pain is nonopioid analgesics. The term "nonopioid analgesics" refer to compounds including, but

10 not limited to Butorphanol, Propoxyphene, meperidine, alphaprodine hydrochloride, fentanyl, and tramadol.

Another preferred group of Drug Used in the Treatment of Pain is "analgesics and antipyretics" wherein the term refers to compounds such as, but not limited to,

15 acetaminophen, ketorolac, allopurinol, methysergide maleate, and methotrimeprazine.

Applicants appreciate that a new Drug Used in the Treatment of Pain may be in development, and the present invention contemplates a synergistic composition comprising

20 such new agents with atypical antipsychotic as well.

As used herein the term "atypical antipsychotic" shall refer to a compound selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine.


25 Risperidone is a known antipsychotic compound currently marketed by Janssen and claimed by U.S. Patent No. 5,246,935 which is hereby incorporated by reference in its entirety.

30 Clozapine is a well known atypical antipsychotic compound currently marketed by Sandoz.


Seroquel is a known compound claimed by U.S. Patent 4,879,288 which is hereby incorporated by reference in its entirety.

35 Sertindole is a known compound and is claimed by U.S. Patent Nos. 5,112,838 and 5,223,945 each of which is hereby incorporated by reference in their entirety.

Ziprasidone is a known compound and is claimed in EP281309-A which is readily available to the skilled artisan. Ziprasidone has the following structure:



5                      Zotepine is a known compound claimed in U.S. Patent No. 3,704,245 which is hereby incorporated by reference in its entirety. Zotepine has the following structure:



10                      As used herein, the term "mammal" shall refer to the Mammalia class of higher vertebrates. The term "mammal" includes, but is not limited to, a human. The term "treating" as used herein includes prophylaxis of the named

condition or amelioration or elimination of the condition once it has been established.

As used herein the term "Drug Used in the Treatment of Pain" refers to compounds known to be clinically useful as analgesics. The term refers to one or more such compounds. Thus, the term Drug Used in the Treatment of Pain can refer to one known analgesic or a combination comprising from two to three known analgesic compounds. Drug Used in the Treatment of Pain are most preferably selected from the compounds named herein.

In the composition of this invention an atypical antipsychotic or a pharmaceutically acceptable salt thereof and one or more Drug Used in the Treatment of Pain are combined in a weight ratio of atypical antipsychotic to Drug Used in the Treatment of Pain of from about one part atypical antipsychotic to from about 1 to about 1000 parts Drug Used in the Treatment of Pain.

A preferred composition is a weight ratio of atypical antipsychotic to another Drug Used in the Treatment of Pain is from about 1 part atypical antipsychotic to from about 1 part Drug Used in the Treatment of Pain to about 100 parts Drug Used in the Treatment of Pain. An especially preferred ratio is from about 1 to about 30. A further preferred ratio may be from about 1 to about 10. A final preferred ratio may be from about 1 to about 3.

Atypical antipsychotics are effective over a wide dosage range; however, it is desirable to administer a dosage that is as low as possible. The amount of Drug Used in the Treatment of Pain present in the composition is adjusted as described above in ratio to the atypical antipsychotic dosage. For example, dosages per day of the atypical antipsychotic will normally fall within the range of about 0.5 mg to about 300 mg per day and the Drug Used in the Treatment of Pain in the composition would be from 3 to 1000 times this amount. However, it will be understood that the amount of the compound actually administered will be determined by a physician, in the light of the relevant



circumstances including the condition to be treated, the choice of compound to be administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration, and therefore the above dosage ranges are not intended to limit the scope of the invention in any way. While the present compounds are preferably administered orally to humans susceptible to or suffering from pain, the compounds may also be administered by a variety of other routes such as the transdermal, parenterally, subcutaneous, intranasal, intramuscular and intravenous routes. Such formulations may be designed to provide delayed or controlled release using formulation techniques which are known in the art.

As used herein the term "treating" includes prophylaxis of a physical and/or mental condition or amelioration or elimination of the developed physical and/or mental condition once it has been established or alleviation of the characteristic symptoms of such condition.

As used herein the term "pain" shall refer to all types of pain. Preferred, the term shall refer to chronic pains, such as neuropathic pain, and post-operative pain, chronic lower back pain, cluster headaches, herpes neuralgia, phantom limb pain, central pain, dental pain, neuropathic pain, another Drug Used in the Treatment of Pain -resistant pain, visceral pain, surgical pain, bone injury pain, pain during labor and delivery, pain resulting from burns, including sunburn, post partum pain, migraine, angina pain, and genitourinary tract-related pain including cystitis, the term shall also preferredly refer to nociceptive pain or nociception.

The dosage administered will, of course, vary depending on known factors such as the pharmacodynamic characteristics of the particular agent, and its mode and route of administration; age, health, and weight of the recipient; nature and extent of the symptoms, kind of concurrent treatment, frequency of treatment, and the effect desired. Usually, the daily dosage can be such

that the active ingredient is administered at a daily dosage of from about 0.2 mg to about 50 mg atypical antipsychotic and from about 0.6 to about 500 mg of another Drug Used in the Treatment of Pain s.

5                   Compositions suitable for internal administration contain from about one half (0.5) milligrams to about 600 milligrams of active ingredient per unit. In these pharmaceutical compositions, the active ingredient will ordinarily be present in an amount  
10 of from about 0.5% to about 95% by weight based on the total weight of the composition.

                  Typical compositions include atypical antipsychotic or a pharmaceutically acceptable acid addition salt thereof and one or more another Drug Used  
15 in the Treatment of Pain s, associated with a pharmaceutically acceptable excipient which may be a carrier, or a diluent or be diluted by a carrier, or enclosed within a carrier which can be in the form of a capsule, sachet, paper, or other container. In making  
20 the compositions, conventional techniques for the preparation of pharmaceutical compositions may be used. For example, the active compound will usually be mixed with a carrier, or diluted by a carrier, or enclosed within a carrier which may be in the form of a ampoule,  
25 capsule, sachet, paper, or other container. When the carrier serves as a diluent, it may be solid, semi-solid, or liquid material which acts as a vehicle, excipient, or medium for the active compound. The active compound can be adsorbed on a granular solid container for example in  
30 a sachet. Some examples of suitable carriers are water, salt solutions, alcohols, polyethylene glycols, polyhydroxyethoxylated castor oil, gelatine, lactose, amylose, magnesium stearate, talc, silicic acid, fatty acid monoglycerides and diglycerides, pentaerythritol  
35 fatty acid esters, hydroxymethylcellulose and polyvinylpyrrolidone. The formulations may also include wetting agents, emulsifying and suspending agents,

preserving agents, sweetening agents, or flavoring agents. The formulations of the invention may be formulated so as to provide quick, sustained, or delayed release of the active ingredient after administration to the patient by employing procedures well known in the art.

The pharmaceutical preparations can be sterilized and mixed, if desired, with auxiliary agents, emulsifiers, salt for influencing osmotic pressure, buffers and/or coloring substances and the like, which do not deleteriously react with the active compounds.

For parenteral application, particularly suitable are injectable solutions or suspensions, preferably aqueous solutions with the active compound dissolved in polyhydroxylated castor oil.

Tablets, dragees, or capsules having talc and/or a carbohydrate carrier or binder or the like are particularly suitable for oral application. Preferable carriers for tablets, dragees, or capsules include lactose, corn starch, and/or potato starch. A syrup or elixir can be used in cases where a sweetened vehicle can be employed.

The compositions of this invention may be suitable for administration to an animal. Such animals include both domestic animals, for example livestock, laboratory animals, and household pets, and non-domestic animals such as wildlife. More preferred, the animal is a vertebrate. Most preferred, a compound of this invention shall be administered to a mammal. It is especially preferred that the animal is a domestic mammal or a human. The most preferred mammal is a human. For such purposes, a compound of this invention may be administered as a feed additive.

#### 35                    Utility Test Methods

The unexpectedly enhanced analgesic activity of the composition of the invention is evidenced by tests

initially conducted on mice. Mice weighing from about 18-25 grams at the time of testing are used for the following studies. All mice are dosed by the oral route with a Drug Used in the Treatment of Pain and/or an atypical antipsychotic.

#### Mouse Writhing Test

An accepted standard for detecting and comparing the analgesic activity of different classes of analgesic compounds for which there is a good correlation with human analgesic activity is the prevention of acetic acid induced writhing in mice. [R. Koster et al. Acetic acid for analgesic screening. Fed. Proc. 18:412, 1959].

Mice, treated with various doses of atypical antipsychotic, another Drug Used in the Treatment of Pain, an atypical antipsychotic: Drug Used in the Treatment of Pain composition, or vehicle are injected intraperitoneally with a standard challenge dose of acetic acid 5 minutes prior to a designated observation period. The acetic acid is prepared as a 0.55% solution and injected at a volume of 0.1 ml/10 grams of body weight. For scoring purposes a "writhe" is indicated by whole body stretching or contracting of the abdomen during an observation period beginning about five minutes after the administration of acetic acid.

#### Sciatic Nerve Ligation Model

An accepted model for assessment of neuropathic pain analgesia is the sciatic nerve ligation model [Bennett, G.J. and Xie, Y.-K. A peripheral mononeuropathy in rat that produces disorders of pain sensation like those seen in man. Pain 33 (1988) 87-107; Lee, Y.-W., Chaplan, S.R. and Yaksh, T.L.: Systemic and supraspinal, but not spinal, opiates suppress allodynia in a rat neuropathic pain model. Neurosci Lett 186 (1995) 111-114]. Rats are anesthetized and a nerve ligation procedure performed. The common sciatic nerve is

exposed and 4 ligatures tied loosely around it with about 1 mm spacing. One day to 10 weeks after surgery, the nociceptive testing is performed. Responses to noxious heat are determined by placing the rats in a chamber with a clear glass floor and aiming at the plantar surface of the affected foot a radiant heat source from beneath the floor. Increased latency to withdraw the hindpaw is demonstrative of analgesic activity. Responses to normally innocuous mechanical stimuli is determined by placing the rats in a chamber with a screen floor and stimulating the plantar surface of the hind paw with graduated von Frey hairs which are calibrated by the grams of force required to bend them. Rats with sciatic nerve ligation respond to lower grams of mechanical stimulation by reflexive withdrawal of the foot than unoperated rats. This response to stimuli which are normally innocuous is termed allodynia. Increases in the grams of mechanical force required to produce foot withdrawal is demonstrative of antiallodynic activity.

#### Formalin Test

The formalin test is a well accepted model of inflammatory pain [Malmberg, A.B. and Yaksh, T.L.: Antinociceptive actions of spinal nonsteroidal anti-inflammatory agents on the formalin test in the rat. The Journal of Pharmacology and Experimental Therapeutics 263 (1992) 136-146]. Rats are anesthetized and when there is a loss of spontaneous movement they are injected subcutaneously in the dorsal surface of the hindpaw with 50 µl of 5% formalin solution using a 30 gauge needle. Rats are then individually placed in an open Plexiglas chamber for observation, and within a maximum interval of 1 to 2 min, the animals display recovery from anesthesia with spontaneous activity and normal motor function. Pain behavior is quantified by periodically counting the incidents of spontaneous flinching/shaking of the injected

paw. The flinches are counted for 1-min periods at 1- to 2-, 5- to 6- and 5min intervals during the interval from 10 to 60 min. Inhibition of the pain behavior is demonstrative of an analgesic activity.

5 All  $ED_{50}$  values and their standard errors of the mean (S.E.M.) are determined using accepted numerical methods. For example, see R. E. Kirk (1982) Experimental Design: Procedures for the behavioral sciences, 2nd ed. Belmont, CA: Brooks/Cole Publishing Co. The interaction of  
10 the dosages on analgesia is demonstrated graphically by the Loewe isobologram (S. Loewe, Pharm. Rev. 9:237-242, 1957).

The interaction of an atypical antipsychotic and another compound used in the treatment of pain on analgesia is demonstrated by Loewe isobologram analysis. In the  
15 isobolographic analysis, the analgesic effects of an atypical antipsychotic are presented on the X-axis and of the other compound used in the treatment of pain on the Y-axis. The line connecting the  $ED_{50}$  dosages of an atypical antipsychotic alone and another compound used in the  
20 treatment of pain alone represents the "ED50 addition line" which indicates the expected location of the  $ED_{50}$  values for an atypical antipsychotic and another compound used in the treatment of pain combinations if simple additivity were to describe their combined effects. According to Loewe's  
25 isobolographic theory, if the analgesic effects of an atypical antipsychotic and another compound used in the treatment of pain were simply additive to one another, the expected location of the  $ED_{50}$  values of the an atypical antipsychotic and another compound used in the treatment of  
30 pain components of each fixed dosage ratio would lie on the addition line. Combination  $ED_{50}$  values located significantly below the  $ED_{50}$  addition line would represent unexpectedly enhanced analgesic activity and combination

-13-

ED<sub>50</sub> values located above the line would represent unexpected diminished analgesic effect.

One method to establish the significance of such unexpected enhanced or diminished activity is to calculate the SEM values for each ED<sub>50</sub>. If the SEM values do not overlap the line of addition, then the ED<sub>50</sub> values are significantly different from the line of addition.

Surprisingly, such experiments demonstrate that compositions comprised of an atypical antipsychotic and another compound used in the treatment of pain show a statistically significant synergistic analgesic effect.

It will be apparent that the instant specifications and examples are set forth by way of illustration and not limitation, and that various modifications and changes may be made without departing from the spirit and scope of the present invention.

Such experiments support that atypical antipsychotics and atypical antipsychotic:another Drug Used in the Treatment of Pain compositions can provide an analgesic effect. Such compositions can provide a statistically significant synergistic analgesic effect.

#### Clinical observations.

A double-blind multicenter clinical trial is designed to assess the safety and efficacy of the atypical antipsychotic. Patients are randomized to atypical antipsychotic, atypical antipsychotic: another Drug Used in the Treatment of Pain composition of this invention, another Drug Used in the Treatment of Pain alone, or placebo. Patients are monitored for perception of pain using standard methods.

The materials for the present invention can be purchased or prepared by a variety of procedures well known

to those of ordinary skill in the art. The atypical antipsychotic compounds are either commercially available or can be prepared using methods described in the patents incorporated herein by reference or as described in widely available publications.

The following examples are provided for purposes of illustration and are not to be construed as limiting the scope of the claimed invention.

#### EXAMPLE 1

A portion of the hydroxypropyl cellulose was dissolved in purified water to form a solution for granulation. The remaining hydroxypropyl cellulose (total of 4.0% w/w final tablet weight), which was an extra fine grade, was combined with the atypical antipsychotic (1.18% w/w), another Drug Used in the Treatment of Pain (3 % w/w), lactose (79.32% w/w) and a portion of the crospovidone (5% w/w) in a high shear granulator. All ingredients were security sieved prior to addition and dry blended in the granulator. This mixture was then granulated with the hydroxypropyl cellulose solution in the high shear granulator. The granulation was wet sized using standard methods. The wet granulation was then dried in a fluidized bed dryer and sized. The material was then added to a tumble bin mixer.

The running powders consisting of microcrystalline cellulose (granular) (10% w/w), magnesium stearate (0.5% w/w), and the remainder of the crospovidone were added to the sized granulation. The mixture was blended and compressed with the appropriate tooling on tablet compression equipment.

#### Subcoating:

Hydroxypropyl methylcellulose (10% w/w) was mixed with purified water to form a solution. Core tablets were



-15-

divided into approximately equal sections and spray coated with the hydroxypropyl methylcellulose solution. The operation was performed in a perforated coating pan.

5     Coating of Core Tablets:

          Color Mixture White (hydroxypropyl methylcellulose, polyethylene glycol, polysorbate 80, and titanium dioxide) was mixed with purified water to form the coating suspension. Subcoated tablets were divided into approximately equal sections and spray coated with the coating suspension described above. The operation was performed in a perforated coating pan.

15     The coated tablets were lightly dusted with carnauba wax and imprinted with appropriate identification.

EXAMPLE 2

20             A portion of the hydroxypropyl cellulose was dissolved in purified water to form a solution for granulation. The remaining hydroxypropyl cellulose (total of 4.0% w/w final tablet weight), which was an extra fine grade, was combined with the atypical antipsychotic (1.18% w/w), lactose (79.32% w/w) and a portion of the crospovidone (5% w/w) in a high shear granulator. All ingredients were security sieved prior to addition and dry blended in the granulator. This mixture was then granulated with the hydroxypropyl cellulose solution in the high shear granulator. The granulation was wet sized using standard methods. The wet granulation was then dried in a fluidized bed dryer and sized. The material was then added to a tumble bin mixer.

35     The running powders consisting of microcrystalline cellulose (granular) (10% w/w), magnesium stearate (0.5% w/w), and the remainder of the crospovidone were added to the sized

-16-

granulation. The mixture was blended and compressed with the appropriate tooling on tablet compression equipment.

Subcoating:

5

Hydroxypropyl methylcellulose (10% w/w) was mixed with purified water to form a solution. Core tablets were divided into approximately equal sections and spray coated with the hydroxypropyl methylcellulose solution. The operation was performed in a perforated coating pan.

10

Coating of Core Tablets:

15

Color Mixture White (hydroxypropyl methylcellulose, polyethylene glycol, polysorbate 80, and titanium dioxide) was mixed with purified water to form the coating suspension. Subcoated tablets were divided into approximately equal sections and spray coated with the coating suspension described above. The operation was performed in a perforated coating pan.

20

The coated tablets were lightly dusted with carnauba wax and imprinted with appropriate identification.

25

## Claims

1. A composition for treating pain comprising  
an analgesic dose of an atypical antipsychotic selected  
5 from the group consisting of risperidone, clozapine,  
seroquel, sertindole, ziprasidone, and zotepine or a  
pharmaceutically acceptable salt or solvate thereof;  
and one or more Drug Used in the Treatment of Pain in a  
weight ratio of atypical antipsychotic to another Drug Used  
10 in the Treatment of Pain from about one part atypical  
antipsychotic to from about one (1) to about one thousand  
(1000) parts Drug Used in the Treatment of Pain.

2. A composition of **Claim 1** wherein the another  
15 Drug Used in the Treatment of Pain is selected from the  
group consisting aspirin, ibuprophen, acetaminophen,  
indomethacin, Tylenol #3, tricyclic antidepressants (for  
example desipramine, imipramine, amitriptyline,  
nortriptyline), anticonvulsants (for example,  
20 carbamazepine, valproate), and serotonin reuptake  
inhibitors (for example, fluoxetine, paroxetine,  
sertraline), mixed serotonin-norepinephrine reuptake  
inhibitors (for example venlafaxine, duloxetine),  
serotonin receptor agonists and antagonists, cholinergic  
25 (muscarinic and nicotinic) analgesics, adrenergic agents,  
and neurokinin antagonists.

3. A composition of **Claim 1** wherein the  
atypical antipsychotic is risperidone.

4. A composition of **Claim 2** wherein the Drug  
Used in the Treatment of Pain is selected from the group  
consisting of acetaminophen, cholinergic analgesics, and  
neurokinin antagonists.

35

-18-

5. A composition of Claim 1 wherein the weight ratio of atypical antipsychotic to another Drug Used in the Treatment of Pain is from about one part atypical antipsychotic to from about one (1) part to from about one hundred (100) parts Drug Used in the Treatment of Pain.

6. A composition of Claim 5 wherein the Drug used in the Treatment of Pain is selected from the group consisting of acetaminophen, meperidine, alphaprodine hydrochloride, fentanyl, tramadol, ketorolac, allopurinol, methysergide maleate, methotrimeprazine, and indomethacin.

7. A composition of Claim 5 wherein the weight ratio of atypical antipsychotic to another Drug Used in the Treatment of Pain is from about one part atypical antipsychotic to from about one (1) part to from about ten (10) parts Drug Used in the Treatment of Pain.

8. A composition of Claim 7 wherein the weight ratio of atypical antipsychotic to another Drug Used in the Treatment of Pain is from about one part atypical antipsychotic to from about one (1) part to from about three (3) parts Drug Used in the Treatment of Pain.

9. A composition of Claim 8 wherein the Drug Used in the Treatment of Pain is acetaminophen.

10. A composition of Claim 1 wherein the atypical antipsychotic is clozapine.

11. A composition of Claim 1 wherein the atypical antipsychotic is seroquel.

35

12. A composition of **Claim 1** wherein the atypical antipsychotic is sertindole.

5 13. A composition of **Claim 1** wherein the atypical antipsychotic is ziprasidone.

14. A composition of **Claim 1** wherein the atypical antipsychotic is zotepine.

10 15. A method for treating pain comprising administering an analgesic dose of a composition comprising an atypical antipsychotic selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine or a  
15 pharmaceutically acceptable salt or solvate thereof; and one or more Drug Used in the Treatment of Pain in a weight ratio of atypical antipsychotic to Drug Used in the Treatment of Pain of from about one (1) part atypical antipsychotic to from about one (1) part to about one  
20 thousand (1000) parts Drug Used in the Treatment of Pain.

16. A method of **Claim 15** wherein the Drug Used in the Treatment of Pain is selected from the group consisting of group consisting of acetaminophen,  
25 indomethacin, Tylenol #3, tricyclic antidepressants (for example desipramine, imipramine, amitriptyline, nortriptyline), anticonvulsants (for example, carbamazepine, valproate), and serotonin reuptake inhibitors (for example, fluoxetine, paroxetine,  
30 sertraline), mixed serotonin-norepinephrine reuptake inhibitors (for example venlafaxine, duloxetine), serotonin receptor agonists and antagonists, cholinergic (muscarinic and nicotinic) analgesics, adrenergic agents, and neurokinin antagonists.

35

17. A method of **Claim 16** wherein Drug Used in the Treatment of Pain is selected from the group consisting of acetaminophen, meperidine, alphaprodine hydrochloride, fentanyl, tramadol, ketorolac, 5 allopurinol, methysergide maleate, methotrimeprazine, and indomethacin.

18. A method for treating pain in a mammal comprising administering an analgesic dose an atypical 10 antipsychotic selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine or a pharmaceutically acceptable salt or solvate thereof; to a mammal in need of such treatment.

15 19. A method of **Claim 18** wherein the atypical antipsychotic is Clozapine.

20 20. A method of **Claim 18** wherein the atypical antipsychotic is risperidone.

21. A method of **Claim 18** wherein the atypical antipsychotic is seroquel.

25 22. A method of **Claim 18** wherein the atypical antipsychotic is sertindole.

23. A method of **Claim 18** wherein the atypical antipsychotic is ziprasidone.

30 24. A method of **Claim 18** wherein the atypical antipsychotic is zotepine.

35 25. A method of **Claim 18** wherein pain is neuropathic pain.

-21-

26. A method of Claim 18 wherein pain is nociceptive pain.

5 27. An atypical antipsychotic selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine or a pharmaceutically acceptable salt or solvate thereof; for use in the treatment of pain.

10 28. An atypical antipsychotic selected from the group consisting of risperidone, clozapine, seroquel, sertindole, ziprasidone, and zotepine or a pharmaceutically acceptable salt or solvate thereof; for use as an analgesic.

15

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/04699

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/55, 31/495, 31/445, 31/16

US CL :514/211, 253, 254, 255, 323, 629

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/211, 253, 254, 255, 323, 629

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched  
noneElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)  
cas-online, aps

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 3,704,245 A (UMIO et al.) 28 November 1972, see the entire document.	1, 5, 7, 8, 14, 15, 18, and 24-28
Y	US 4,879,288 (WARAWA et al.) 07 November 1989, see the entire document.	1, 5, 7, 8, 11, 15, 18, 21 and 25-28
Y	US 5,045,539 A (HELSLEY et al.) 03 September 1991, see the entire document.	1, 5, 7, 8, 10, 15, 18, 19 and 25-28
Y	US 5,112,838 A (PERREGAARD et al.) 12 May 1992, see the entire document.	1, 5, 7, 8, 12, 15, 18, 22 and 25-28

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:			
*A	document defining the general state of the art which is not considered to be of particular relevance	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E	earlier document published on or after the international filing date	*X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L	document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O	document referring to an oral disclosure, use, exhibition or other means		
*P	document published prior to the international filing date but later than the priority date claimed	*G	document member of the same patent family

Date of the actual completion of the international search

04 JUNE 1997

Date of mailing of the international search report

09.07.97

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Authorized officer

KEVIN E. WEDDINGTON

Facsimile No. (703) 305-3230

Telephone No. (703) 308-1235



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US97/04699

## C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5,246,935 A (JEPPESEN et al.) 21 September 1993, see the entire document.	1, 3, 5, 7, 8, 15, 18, 20 and 25-28
Y	DAHLIN et al., "THE MERCK INDEX, AN ENCYCLOPEDIA OF CHEMICALS, DRUGS, AND BIOLOGICALS" published 1983 by Merck & Co., Inc. (N.J.), see page 7, abstract No. 39.	2, 4, 6, 9 and 16





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>A61K 31/55, 31/495, 31/445, 31/16</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 97/35584</b> <b>(43) International Publication Date:</b> 2 October 1997 (02.10.97)
<b>(21) International Application Number:</b> PCT/US97/04699 <b>(22) International Filing Date:</b> 24 March 1997 (24.03.97)  <b>(30) Priority Data:</b> 60/014,152      25 March 1996 (25.03.96)      US  <b>(71) Applicant:</b> ELI LILLY AND COMPANY [US/US]; Lilly Corporate Center, Indianapolis, IN 46285 (US).  <b>(72) Inventors:</b> HELTON, David, R.; 2607 South Bo-Mar Lane, Greenfield, IN 46140 (US). SHANNON, Harlan, E.; 4229 Rolling Springs Drive, Carmel, IN 46234 (US). WOMER, Daniel, E.; Apartment H., 2121 Blue Jay Court, Indianapolis, IN 46260 (US).  <b>(74) Agents:</b> VORNDRAN-JONES, MaCharri et al.; Eli Lilly and Company, Lilly Corporate Center, Indianapolis, IN 46285 (US).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NQ, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> METHOD FOR TREATING PAIN  <b>(57) Abstract</b>  The present invention provides a method for treating pain using an atypical antipsychotic compound.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NI	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						